

The Outer Regions of the Nearby Sc Galaxies NGC 2403 and M33: Evidence for an Intermediate Age Population at Large Radii ¹

T. J. Davidge

Canadian Gemini Office, Herzberg Institute of Astrophysics,
National Research Council of Canada, 5071 West Saanich Road,
Victoria, B.C. Canada V9E 2E7
email: tim.davidge@nrc-cnrc.gc.ca

ABSTRACT

Deep $g'r'i'z'$ images obtained with the Gemini Multi-Object Spectrograph (GMOS) on Gemini North are used to investigate the stellar content in the outer regions of the nearby Sc galaxies NGC 2403 and M33. The field observed in NGC 2403 covers galactocentric distances between 5 and 11 kpc perpendicular to the line of sight (R_{GC}^{LOS}), and 7 and 19 kpc along the plane of the disk (R_{GC}^{disk}). The red giant branch (RGB)-tip occurs at $i' = 23.6 \pm 0.1$, and the Cepheid and RGB-tip distance scales for NGC 2403 are in good agreement. The number density of bright main sequence stars in this field experiences a steep cut-off at $R_{GC}^{disk} \sim 10$ kpc, which is consistent with the expected truncation radius of the disk predicted from studies of edge-on spiral galaxies. While very young stars are restricted to $R_{GC}^{disk} < 10$ kpc, a population of bright asymptotic giant branch (AGB) stars is present throughout the entire GMOS field, indicating that star formation occurred outside of the present-day star-forming disk of NGC 2403 during intermediate epochs. The AGB stars are not in a tidal stream; in fact, the ratio of AGB stars above the RGB-tip to those below the RGB-tip does not change with radius, indicating that the bright AGB stars are uniformly mixed with the fainter stellar content throughout the field. The AGB luminosity function (LF) scales with r -band surface brightness over a wide range of radii throughout the main body of NGC 2403, indicating that the age distribution of stars in the outer regions of the present-day star-forming disk is not skewed to

¹Based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a co-operative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Particle Physics and Astronomy Research Council (United Kingdom), the National Research Council of Canada (Canada), CONICYT (Chile), the Australian Research Council (Australia), CNPq (Brazil), and CONICET (Argentina).

younger values than in the inner disk. Based on the color of stars on the upper portions of the RGB it is concluded that metallicity changes across the field, with $[\text{Fe}/\text{H}] = -0.8 \pm 0.1$ (random) ± 0.3 (systematic) at $R_{GC}^{LOS} = 5$ kpc, and $[\text{Fe}/\text{H}] = -2.2 \pm 0.2$ (random) ± 0.8 (systematic) at $R_{GC}^{LOS} = 11$ kpc.

The M33 field samples R_{GC}^{LOS} between 8 and 10 kpc and R_{GC}^{disk} between 14 and 17 kpc. Bright AGB stars are detected in this field, and the ratio of bright AGB stars to stars on the upper RGB is at least as large as that measured in the outer regions of NGC 2403; thus, an intermediate-age population occurs well outside of the young star-forming disk of M33. The color of stars on the upper RGB suggests that $[\text{Fe}/\text{H}] = -1.0 \pm 0.3$ (random) ± 0.3 (systematic) in this field.

The globular cluster systems of both NGC 2403 and M33 contain objects that formed during intermediate epochs, and it is suggested that the luminous AGB stars detected in the current study are the field counterparts of these clusters. The detection of intermediate-age stars in the outer regions of these galaxies is consistent with models in which late-type spiral galaxies formed more slowly than earlier-type systems.

Subject headings: galaxies: individual (NGC 2403, M33) - galaxies: spiral - galaxies: halo - galaxies: stellar content - galaxies: evolution - stars: AGB and post-AGB

1. INTRODUCTION

The halos and outer disks of spiral galaxies are basic laboratories for studying the formation and evolution of these systems. Numerical simulations predict that halo formation may be an extended, and perhaps even on-going, process that involves the merging of proto-systems having a range of masses and chemical enrichment histories (e.g. Bekki & Chiba 2001). The observational evidence that the outer regions of the Milky-Way contains stars accreted from companion galaxies is now well established. In particular, there are a handful of globular clusters in the Milky-Way that are markedly younger than the majority (e.g. Rosenberg et al. 1999; Buonanno et al. 1998; Sarajedini, Chaboyer, & Demarque 1997), as well as tidal streams containing stars that are distinct from those in the present day halo field (e.g. Ibata et al. 2001a; Martinez-Delgado et al. 2001; Helmi & White 1999; Majewski, Munn, & Hawley 1994); similar structures are also seen in the outer regions of M31 (Ibata et al. 2001b; Saito & Iye 2000). The overall structural characteristics

of the Galactic halo are also consistent with formation driven by hierarchal accretion (e.g. Chiba & Beers 2001; Bullock, Kravtsov, & Weinberg 2001), although there remain difficulties explaining the presence of a thin disk, given that interactions with satellites heat, and thereby thicken, the disk (Chiba & Beers 2001).

It is not clear if the age and metallicity distribution function (MDF) of halo stars depend on global host galaxy properties such as mass and morphology. The stochastic processes inherent to hierarchal evolution (e.g. Côté et al. 2000), including differences in the gas fraction of accreted satellites (Harris & Harris 2001), are expected to smear any such relations. Studies of the brightest resolved stars in the halos of Local Group galaxies have revealed intriguing galaxy-to-galaxy differences, although the number of systems studied to date is too small to allow secure trends to be defined. The halos of M31, the Milky-Way, and M33 have very different mean metallicities (e.g. Mould & Kristian 1986 - hereafter MK86; Durrell, Harris, & Pritchett 1994, 2001), in the sense that the M31 halo is the most metal-rich of the three. The globular cluster systems of these galaxies also show differences. The depths of CN absorption bands in the integrated spectra of M31 and Milky-Way clusters are suggestive of different chemical mixtures (Burstein et al. 1984, Davidge 1990), while M33 contains a significant number of intermediate age globular clusters (Christian & Schommer 1988 – hereafter CS88; Sarajedini et al. 1998; Chandar et al. 2002) that do not have counterparts in either M31 or the Milky-Way. With HST the investigation of resolved stars in halos has been expanded to galaxies outside the Local Group. Harris & Harris (2000; 2001) point out that the halo MDF of the elliptical galaxy NGC 5128, which is more massive than M31, is similar to the MDFs of systems that are as diverse as M32 (Grillmair et al. 1996), the LMC (Cole, Smecker-Hane, & Gallagher 2000), and the M31 halo (Durrell et al. 2001), which in turn suggests of similarities in their early evolution.

Disks are thought to form from material that was subjected to tidal torques during early epochs (e.g. Zurek, Quinn, & Salmon 1988). Feedback from star formation, coupled with interactions between the host galaxy and satellites on plunging orbits, will heat proto-disk material and thereby delay disk formation (e.g. Weil, Eke, & Efstathiou 1998). In the case of the Milky-Way, the oldest stars in the Galactic disk have ages that are consistently lower than those inferred for typical globular clusters (e.g. Oswalt et al. 1996; Leggett, Ruiz, & Bergeron 1998; Knox, Hawkins, & Hambly 1999; Liu & Chaboyer 2000, but see also Binney, Dehnen, & Bertelli 2000), which is consistent with a hiatus between the formation of the halo and the disk. In fact, the youngest globular clusters have ages that are similar to the oldest disk stars, suggesting that the formation of the Galactic disk commenced only after the conditions required to form globular clusters were suppressed.

Disks might be expected to form from the inside out, as the low densities of the

outer disk make these regions more fragile than the inner disk, and hence more easily disrupted by disturbances from nearby satellites. The outer disk may also experience delayed star formation if infall and the viscous transport of material from smaller radii have comparable time scales (e.g. Ferguson & Clarke 2001). The inside-out disk formation model is supported by the presence of age gradients in the disks of nearby galaxies, in the sense that mean age decreases (becomes younger) towards larger radii (e.g. Bell & de Jong 2000). There is also a tendency for the region near the disk boundary in edge-on systems to be bluer than in the rest of the disk, as expected if the disk/halo interface is dominated by young stars, although there are some exceptions (de Grijs, Kregel, & Wesson 2001).

Deep photometric studies of the resolved stellar contents of nearby spiral galaxies offer a direct means of probing the evolution of the outer regions of these systems. In this paper deep $g'r'i'z'$ images obtained with the Gemini Multi-Object Spectrograph (GMOS) on Gemini North are used to probe the bright stellar content in the outer regions of the nearby Sc galaxies M33 and NGC 2403. The fields observed in these galaxies have similar galactocentric radii, R_{GC} , thereby permitting a direct comparison of stellar contents.

NGC 2403 and M33 have roughly the same M_B , morphology, inclination, gas-phase metal abundances (Garnett et al. 1997), and star-forming histories in the inner disk (Davidge & Courteau 2002); hence, they are well matched for a comparative study. These galaxies also share a common peculiarity, in that both are experiencing vigorous star formation in the inner disk, even though the gas densities in these same regions appear to be too low to trigger the collapse instabilities required for star formation (Kennicutt 1989; Martin & Kennicutt 2001). These similarities aside, M33 and NGC 2403 are located in slightly different environments. While both galaxies are in a hierarchal system with a larger Sb galaxy, NGC 2403 is ~ 0.9 Mpc from M81 (Karachentsev et al. 2000), while M33 is only ~ 0.2 Mpc from M31. Furthermore, the M81 group contains dwarf galaxies that do not have known Local Group counterparts (Caldwell et al 1998), and this might affect evolution driven by satellite accretion.

The distance moduli and reddenings adopted for M33 and NGC 2403 throughout this study are listed in Table 1. The distance modulus for NGC 2403 in Table 1 is based on Cepheids, and in §4 it is demonstrated that the Cepheid and RGB-tip distances for NGC 2403 are in good agreement. As for M33, Cepheids and the RGB-tip give distance moduli that differ by 0.3 dex (e.g. Lee et al. 2002), and the RGB-tip distance measured by Kim et al. (2002) has been adopted for this study as it is likely less susceptible to internal reddening.

The observations and data reduction techniques are discussed in §2, while details of the photometric measurements are presented in §3. The NGC 2403 field samples a range of

environments, and in §4, 5, and 6 the stellar contents of three distinct regions, that sample the outer star-forming disk, the disk/halo interface, and the halo, are discussed. The stellar content of the M33 field, which samples a region outside the active star-forming disk, is investigated in §7. A discussion and summary of the results follows in §8.

2. OBSERVATIONS & REDUCTIONS

The data were recorded during Semester 2001B as part of the System Verification program for the GMOS (Crampton et al. 2002) on Gemini North. g' , r' , i' , and z' images were obtained of fields along the minor axes of NGC 2403 and M33; control fields were also observed to monitor contamination from foreground stars and background galaxies. The right ascensions and declinations of the various fields are listed in Table 2, while the locations of the NGC 2403 and M33 fields on the Palomar Sky Survey are shown in Figures 1 and 2. The M33 and NGC 2403 fields have the same central R_{GC} , although a smaller portion of M33 is sampled than NGC 2403 because of the difference in distance. In fact, the distance to NGC 2403 is such that the GMOS field samples a swath from the star-forming disk to the halo of the galaxy.

The data were recorded using a 6 point dither pattern so that the gaps between the individual CCDs that make up the GMOS detector mosaic could be filled when constructing the final images. Each CCD output was binned 2×2 during read-out, so each superpixel covers 0.145 arcsec on a side. Exposure times were selected using the GMOS Integration Time Calculator (ITC) to allow stars on the upper RGB in each galaxy to be detected at the 5 to 10σ level during median conditions. In general, the delivered photometric performance is consistent with the ITC predictions. Various details of the observations, including exposure time and delivered image quality, are listed in Table 2. The exposure times of the NGC 2403 background field g' observations are much shorter than originally intended as the seeing suddenly degraded on the night these data were recorded, making it necessary to stop prematurely these observations.

The data were reduced with tasks in the Gemini IRAF package, using bias frames and twilight sky flats that were recorded at the beginning and/or end of each night. The data reduction sequence consisted of (1) balancing CCD-to-CCD gain differences, (2) subtracting the bias pattern and DC bias level, (3) dividing by twilight sky flat-field frames, and (4) mosaicking the bias-subtracted and flat-fielded outputs from the individual CCD amplifiers to create a single image. Interference fringes were removed from the i' and z' data by subtracting fringe frames that were constructed by combining flat-fielded images of various fields recorded through these filters. The final images for photometric analysis were

constructed by spatially aligning the individual images for each field, median-combining the results, and then trimming to the region of common sky coverage.

The final i' images of each galaxy and the corresponding control field are shown in Figures 3 (NGC 2403) and 4 (M33). The diffuse object near the eastern edge of the NGC 2403 control field is the dwarf irregular galaxy Kar 50, and the photometric properties of resolved stars in this system were the subject of a separate paper (Davidge 2002).

3. PHOTOMETRIC MEASUREMENTS & ANALYSIS STRATEGY

The photometric measurements were made with the point-spread function (PSF) fitting routine ALLSTAR (Stetson & Harris 1988). PSFs were constructed by combining approximately 50 bright stars per image with the DAOPHOT (Stetson 1987) PSF task. Because the PSF stars are bright and relatively isolated, they were also used to determine aperture corrections, which were measured after subtracting neighboring sources. The photometric measurements are in the Vega-based $g'r'i'z'$ system described by Fukugita et al. (1996), and the calibration was set using standard star observations that were obtained during each observing run. The estimated uncertainty in the calibration is ± 0.05 mag, and this is borne out by the reasonably good agreement with fiducial sequences on the $(g' - r', r' - i')$ two-color diagrams (TCDs).

Artificial star experiments were run to assess completeness, estimate photometric uncertainties, and establish a simple means of identifying obviously extended objects. This latter step was done by rejecting objects having a DAOPHOT ‘sharp’ parameter that differed at the $3 - \sigma$ or higher level from that predicted for point sources. The artificial stars were assigned colors that are typical of real stars in the data, and the completeness fractions in g' and i' for each field are shown in Figure 5.

The NGC 2403 GMOS field contains a mix of populations. Young disk stars dominate in the lower right hand corner of Figure 1, while older stars dominate in the upper left hand corner. Given this gradient in stellar content, the NGC 2403 field was divided into three regions, which are indicated in Figure 3. The boundary between Regions I and II roughly tracks the drop-off in the bright blue main sequence component. Deep imaging studies of edge-on spiral galaxies indicate that disks terminate over a ~ 1 kpc interval (e.g. van der Kruit & Searle 1981; de Grijs et al. 2001), and Region II probes the stellar content in a 1 kpc interval beyond the outer edge of Region I. Region III occupies the remainder of the GMOS field. The HI disk of NGC 2403 extends into Regions II and III (Fraternali et al. 2002).

The control field was similarly divided so that identical areas are covered when making comparisons between the galaxy and control datasets. For the remainder of the paper regions in the galaxy fields are designated with the subscript ‘g’, while regions in the control fields are indicated with the subscript ‘c’.

4. NGC 2403 REGION I_g: THE OUTER STAR-FORMING DISK

4.1. The LFs, TCDs, and CMDs of NGC 2403 Region I_g

The g' and i' luminosity functions (LFs) of Region I_g, corrected for incompleteness and contamination from sources in the corresponding portion of the control field, are shown in Figure 6. There is a large number of objects belonging to NGC 2403 in Region I_g, with the onset occurring near $g' = 23.5$ and $i' = 22$. The majority of sources with $i' > 23.5$ are evolving on the RGB (§4.3).

The majority of sources in Regions I_g and I_c have stellar spectral energy distributions (SEDs), and this is demonstrated in Figures 7 and 8, where the $(g' - r', r' - i')$ and $(r' - i', i' - z')$ TCDs of sources in these regions are compared. The stellar sequence from Figure 3 of Schneider et al. (2001, hereafter SDSS) and the $\log(g) = 2.5$ and 4.5 solar metallicity models from Tables 9 and 4 of Lenz et al. (1998) are also shown in these figures. While the majority of sources follow the SDSS sequence on both TCDs when $r' - i' < 0.5$, there are stars in Region I_g with $r' - i' > 0.5$ and $g' - r' > 1.5$ that fall well off of the SDSS sequence on the $(g' - r', r' - i')$ TCD, and stars with similar photometric properties do not occur in Region I_c. We suggest that the stars with $g' - r' > 1.5$ are red giants that are reddened by dust belonging to NGC 2403. That the stars with $g' - r' > 1.5$ form a more-or-less continuous sequence indicates that the stellar content of Region I_g is not subject to uniform levels of internal extinction; rather, there is differential extinction ranging up to $A_{g'} \sim 1.0$ mag, which corresponds to $A_V \sim 0.9$ and $E(B - V) \sim 0.3$. It is also clear from Figure 7 that not all giants in Region I_g are affected by large amounts of internal extinction. The level of internal extinction predicted from the $(g' - r', r' - i')$ TCD is consistent with the range calculated from the Balmer decrement in NGC 2403 HII regions (McCall, Rybski, & Shields 1985; Petersen & Gammelgaard 1996); for comparison, the statistical prescription of Tully & Fouqué (1985) predicts a mean internal extinction of $A_B = 0.34$ mag ($A_V = 0.26$ mag) for NGC 2403 (Pierce & Tully 1992).

The Region I_g and I_c data tend to have $i' - z'$ colors that are smaller at a fixed $r' - i'$ than predicted from the SDSS sequence. In the case of Region I_g, the offset with respect to the SDSS data is in the range 0.05 – 0.10 mag along the $i' - z'$ axis when $r' - i' < 1$, and so

is within the expected uncertainty in the photometric calibration. However, when $r' - i' > 1$ there is a population of objects that are offset from the SDSS sequence by $0.2 - 0.3$ mag in $i' - z'$, and hence appear to have non-stellar SEDs. This part of the $(r' - i', i' - z')$ TCD is populated by high redshift QSOs (Fan 1999), although the density of these objects on the sky is too low to produce the number of sources found here (Anderson et al. 2001).

The $(r', g' - r')$, $(i', r' - i')$, and $(z', i' - z')$ CMDs of Regions I_g and I_c are plotted in Figures 9, 10, and 11. The faint limits of the Region I_g and I_c data differ because of differences in image quality (e.g. Figure 5). Nevertheless, there is a clear population of stars belonging to NGC 2403 in each of the Region I_g CMDs.

4.2. Young Stars in NGC 2403 Region I_g

Region I_g contains a population of sources with $g' - r'$ and $r' - i' < 0$, and the location of these objects on the TCDs indicate that they have stellar SEDs. The solid and dashed lines in the left hand panel of Figure 9 show the main sequence defined by OB stars in the Milky-Way and LMC, translated to match the distance to NGC 2403. The main sequence relation was obtained by transforming the relation between M_V and effective temperature listed by Humphreys & McElroy (1984) into the SDSS photometric system using the relations between (1) r' and V given by Krisciunas, Margon, and Szkody (1998) and (2) $g' - r'$ and effective temperature predicted by the solar metallicity $\log(g) = 4.5$ Lenz et al. (1998) models. The dashed line shows the main sequence reddened only by line-of-sight extinction from Schlegel et al. (1998), while the solid line shows the main sequence reddened by the foreground and a uniform sheet of internal extinction with $E(B - V) = 0.3$, which the distribution of stars on the $(g' - r', r' - i')$ TCD suggests is the maximum internal extinction for red giants in Region I_g (see above).

The amount of reddening in a galaxy is population dependent, in the sense that it is higher for young OB stars than for red giants, since stars diffuse away from dusty star-forming regions as they age (e.g. Zaritsky 1999). The predicted location of the main sequence with only foreground reddening lies at the left hand edge of the blue stellar content in the Region I_g CMD, indicating that not all of the main sequence stars in Region I_g are subject to significant internal extinction. The predicted main sequence with internal reddening of $A_{g'} = 1$ mag passes through a group of stars with r' between 24 and 23, and if these are actual main sequence stars then they have intrinsic brightnesses comparable to the brightest main sequence stars in the Galaxy and the LMC.

The brightest blue and red stars in Region I_g are likely supergiants. To demonstrate

this point, the locus defined by type Ia supergiants in the Galaxy and Magellanic Clouds, based on the M_V and effective temperatures listed in Humphreys & McElroy (1984) and transformed into the SDSS photometric system using the procedure described above for the main sequence relation, is also shown on Figure 9 for the two reddening situations described above. The Ia supergiant sequence in Figure 9 with an internal extinction of $A_{g'} = 1$ mag defines the upper envelope of blue stars in Region I_g , with four stars falling along this sequence.

The presence of bright supergiants and main sequence stars indicates that Region I_g is a site of recent star formation; Region I_g is thus part of the active star-forming disk of NGC 2403.

4.3. RGB Stars in NGC 2403 Region I_g

The RGB-tip in moderately metal-poor globular clusters occurs near $M_{i'} \sim -4$ and $M_{r'} \sim -2.5$ (Davidge et al. 2002), which corresponds to $i' \sim 23.5$ and $r' \sim 25$ at the distance of NGC 2403 with no internal reddening, and $i' \sim 24.0$ and $r' \sim 25.7$ with internal reddening of $A_{g'} = 1$ mag; the RGB-tip thus falls near the faint limits of the $(r', g' - r')$ and $(z', i' - z')$ CMDs. However, a large population of red stars is seen in the $(i', r' - i')$ CMD of Region I_g when $i' > 23.5$.

The i' LF of sources with $r' - i'$ between 0.2 and 0.7 in Region I_g , which is the color range containing the greatest density of RGB stars, is shown in Figure 12. The rate of evolution on the RGB is insensitive to metallicity (e.g. VandenBerg 1992), and so the power-law exponent that characterizes the RGB LF does not vary greatly with metallicity. The LF of RGB stars in globular clusters follows a power law with $\Delta \log(n)/\Delta(i') \sim 0.4$ when $[\text{Fe}/\text{H}] < -1.6$ (Davidge et al. 2002), and a least squares fit to the LF entries when i' is between 23.6 and 25.0 indicates that the exponent is 0.60 ± 0.07 , which is higher than that measured in metal-poor globular clusters. The fitted relation is compared with the observed LF in Figure 12, and there is a tendency for the fitted relation to fall above the LF entries when $i' > 23.6$, indicating that the RGB-tip brightness in NGC 2403 is $i' = 23.6 \pm 0.1$; to the best of our knowledge, this is the first detection of the RGB-tip in NGC 2403.

The Cepheid and RGB-tip distance moduli of M33 differ by 0.3 dex, presumably due to internal extinction (Lee et al. 2002); is this also the case for NGC 2403? Using the Cepheid distance modulus for NGC 2403 listed in Table 1, and assuming that the majority of RGB-tip stars in Region I_g are subject to foreground reddening, then the intrinsic brightness of the RGB-tip in Region I_g is $M_{i'}^{\text{RGBT}} = (23.6 \pm 0.1) - (27.51 \pm 0.24) - (2.086 \times 0.046) = -4.0 \pm 0.3$;

the last term is the correction for extinction due to foreground dust using $\frac{A_{i'}}{E(B-V)} = 2.086$ from Table 6 of Schlegel et al. (1998). This value of $M_{i'}^{RGBT}$ is consistent with that found in metal-poor globular clusters and the dwarf spheroidal galaxy And V by Davidge et al. (2002). While the RGB stars in Region I_g are moderately metal-rich (§8), the RGB-tip brightness in the near-infrared does not change markedly with metallicity when $[\text{Fe}/\text{H}] < -0.7$ (Da Costa & Armandroff 1990). Thus, the Cepheid and RGB-tip distances for NGC 2403 are consistent.

The mean color and width of the RGB provides information about the chemical composition of old and intermediate age populations. In the current study, we use $r' - i'$ to probe RGB color, as the data in r' and i' go much deeper than the RGB-tip, sample red populations (e.g. Figure 10), and are less susceptible to line blanketing than data recorded at shorter wavelengths (e.g. Bica, Barbuy, & Ortolani 1991). The histogram distribution of $r' - i'$ colors of sources in Region I_g with i' between 24 and 25, corrected for sources in the same brightness interval in Region I_c, is shown in the top panel of Figure 13. The analysis is restricted to stars with i' between 24.0 and 25.0 as the r' and i' data are 100% complete at these brightnesses over the range of RGB colors.

The $r' - i'$ distribution for Region I_g has a broad peak, due to the RGB, with a secondary peak when $r' - i' < 0$, which is due to bright main sequence stars (§4.2). The distribution of $r' - i'$ colors about the red peak is symmetric, and the mean, based on the data in bins between 0.0 and 1.0, is $\overline{r' - i'} = 0.50 \pm 0.01$, where the uncertainty is the standard error in the mean. The width of the $r' - i'$ distribution is dominated by photometric errors. To demonstrate this point, the gaussian error distribution predicted from the artificial star experiments, scaled to match the number of stars between $r' - i' = 0.3$ and 0.7 , is compared with the observed distribution in Figure 13, and there is excellent agreement with the observations. Differential reddening of size $\Delta A_{g'} = 1$ mag does not contribute significantly to the width of the observed color distribution. If the standard deviation of the internal extinction is one half of the total internal extinction (i.e. $\sigma_{A'_{g'}} = \pm 0.5$ mag), then the dispersion along the $r' - i'$ axis due to differential reddening is $\sigma_{r'-i'} = \pm 0.1$. The result of adding this component in quadrature to that due to photometric errors is to broaden the total predicted color distribution by only an additional ± 0.02 mag.

4.4. AGB Stars in NGC 2403 Region I_g

AGB stars are seen above the RGB-tip in the Region I_g CMDs. Stars near the AGB-tip have low temperatures, and line blanketing can suppress the brightnesses of these objects at visible wavelengths, causing the AGB to be a near horizontal sequence on CMDs.

The effects of line blanketing become smaller towards longer wavelengths, and the AGB is a vertical sequence on the $(z', i' - z')$ CMD of Region I_g . Hence, the z' LF of stars above the RGB-tip is an obvious direct probe of AGB content.

The z' LFs of sources with $i' - z'$ between 0 and 1 in Region I_g is shown in Figure 14. The RGB-tip occurs near $z' = 23.4$, and the onset of the RGB causes the Region I_c LF to climb when $z' > 23.4$. The LF is remarkably flat between $z' = 22.2$ and 23.4, and there is an apparent discontinuity at $z' = 22$, where the number counts drop.

The presence of bright AGB stars above the RGB-tip suggests that Region I_g experienced star formation during intermediate epochs. To check this result, and also enable comparisons with the AGB contents of other areas of NGC 2403 and other galaxies, the bolometric LF of AGB stars was constructed using brightnesses and colors from the $(i', r' - i')$ CMD of Regions I_g and I_c . Hudon et al. (1989) used V, R , and I images to investigate the AGB content of three fields in NGC 2403 that lie roughly along the major axis of the galaxy. For consistency, the criteria for defining AGB stars and rejecting red supergiants discussed by Hudon et al. (1989) have been adopted for the current study. $r' - i'$ colors and i' magnitudes were transformed into $R - I$ and I (Kron-Cousins system) using relations from Fukugita et al. (1996) after adopting the relation between $V - I$ and $R - I$ for giants from Bessell (1979). Bolometric corrections were computed using Equation 2 of Bessell & Wood (1984).

The AGB LF of Region I_g is shown in Figure 15. The dashed line is the composite AGB LF of Fields 2 and 3 from Hudon et al. (1989), which are located 360 and 480 arcsec from the galaxy center. The Hudon et al. (1989) LFs in Figure 15 have been (1) scaled to match the number of AGB stars in Region I_g with M_{bol} between -4.5 and -6.2 , and (2) shifted by 0.1 mag along the horizontal axis to match the distance modulus adopted here. The dashed-dotted line is the AGB LF of the LMC constructed by Reid & Mould (1984), shifted along the vertical axis to match the number of sources in Region I_g with M_{bol} between -4.5 and -6.2 . Both the Hudon et al. (1989) and Reid & Mould (1984) LFs match the Region I_g LF, suggesting that the star forming histories of Hudon et al. (1989) Fields 2 and 3, the LMC, and Region I_g have been similar during the past few Gyr.

5. NGC 2403 REGION II_g : THE EDGE OF THE YOUNG DISK

5.1. The LFs, TCDs, and CMDs of NGC 2403 Region II_g

The boundary between Regions I and II tracks the drop-off in the number of bright blue sources, and so Region II_g does not contain objects that are as bright as those in

Region I_g . This is evident in the g' and i' LFs of Region II_g , corrected for incompleteness and sources in the corresponding portion of the control field, which are shown in Figure 6. The onset of stars in Region II_g occurs near $i' = 23$.

The $(g' - r', r' - i')$ and $(r' - i', i' - z')$ TCDs of Regions II_g and II_b are shown in the right hand panels of Figures 7 and 8. Whereas the $(g' - r', r' - i')$ TCD of Region I_g has an excess population of objects with respect to Region I_c over a broad range of colors, the $(g' - r', r' - i')$ TCDs of Regions II_g and II_c differ mainly when $g' - r' > 0.6$ and $r' - i' > 0.1$. In fact, the $(g' - r', r' - i')$ TCD of Region II_g contains only two objects with $g' - r' < 0$ that have stellar SEDs, indicating that the number density of bright main sequence stars in Region II_g is lower than in Region I_g . An absence of sources with $r' - i' < 0$ is also evident in the $(r' - i', i' - z')$ TCD of Region II_g .

As was the case in Region I_g , the $(g' - r', r' - i')$ TCD of Region II_g contains sources with $g' - r' > 1.5$ and $r' - i' > 0.4$ that fall well off of the SDSS sequence, whereas Region II_c contains only a modest number of objects with similar colors. The presence of objects with $g' - r' > 1.5$ suggests that, as in Region I_g , some red giants in Region II_g are reddened by dust belonging to NGC 2403.

The $(r', g' - r')$, $(i', r' - i')$, and $(z', i' - z')$ CMDs of Regions II_g and II_c are shown in Figures 9, 10, and 11. While some objects that may be young main sequence stars are present in the $(r', g' - r')$ CMD of Region II_g , the brightest of these have $r' > 24$, and hence are 1 mag fainter than the brightest main sequence stars in Region I_g . While there is a star in the $(r', g' - r')$ CMD of Region II_g that has a brightness and color consistent with it being a Type Ia supergiant, the number density of bright objects in the Region II_c CMD suggests that this may be a foreground star with fortuitous photometric properties.

5.2. RGB Stars in NGC 2403 Region II_g

In §4.3 it was demonstrated that the LF of RGB stars in Region I_g follows a power-law with an exponent that is larger than that in globular clusters. The i' LF of stars in Regions II_g with $r' - i'$ between 0.2 and 0.7, corrected for the number of sources in the same color range in Region II_c , is shown in Figure 12. A least squares fit to sources with i' between 23.6 and 25.0 in the Region II_g LF indicates that the power-law exponent in Region II_g is $\sim 0.27 \pm 0.12$, which is smaller than that in Region I_g , and is not significantly different from that in globular clusters.

The RGBs of Regions I_g and II_g have different mean colors. The $r' - i'$ color distributions of sources with i' between 24 and 25 in Region II_g , corrected for sources in

Region II_c, is shown in Figure 13. The RGB of Region II_g is skewed towards lower $r' - i'$ values than in Region I_g, with $\overline{r' - i'} = 0.42 \pm 0.01$. Metallicity and extinction are two possible causes of the difference between the mean $r' - i'$ colors of upper RGB stars in Regions I_g and II_g². While the presence of sources with $g' - r' > 1.5$ on the $(g' - r', r' - i')$ TCDs of Regions I_g and II_g indicate that both regions contain stars that are subject to reddening internal to NGC 2403, Region I_g contains sources that are redder than in Region II_g, suggesting that the maximum reddenings in the two fields may differ, which would cause $\overline{r' - i'}$ in Region I_g to be larger than in Region II_g. The difference in mean giant branch color between Region I_g and II_g is $\Delta(r' - i') = 0.08 \pm 0.02$ and, if due entirely to reddening, this corresponds to $\Delta A_V = 0.4 \pm 0.1$ mag, which is consistent with the difference between the $g' - r'$ colors of the reddest sources in the $(g' - r', r' - i')$ TCDs of Regions I_g and II_g. However, it is evident from the TCDs that not all RGB stars in Regions I_g and II_g are subject to this high level of extinction. Moreover, in §6.2 it is demonstrated that $\overline{r' - i'}$ in Region III_g is lower than in Regions I_g and II_g by an amount that is too large to be explained by extinction. Extinction is therefore not a convincing explanation for the difference in RGB color seen throughout the GMOS field.

5.3. AGB Stars in NGC 2403 Region II_g

AGB stars form a spray of objects above the RGB-tip in the Region II_g $(i', r' - i')$ and $(z', i' - z')$ CMDs, and the presence of these stars indicates that Region II_g experienced star formation during intermediate epochs. The overall morphologies of the $(z', i' - z')$ CMDs of Regions I_g and II_g are similar: as in Region I_g, the AGB sequence in the $(z', i' - z')$ CMD of Region II_g is almost vertical, and peaks near $z' \sim 22$. These similarities suggest that Regions I_g and II_g may have similar AGB contents.

The z' LF of Region II_g is shown in Figure 14. The z' LFs of Regions II_g and I_g are similar, in that both are more-or-less flat in the interval between the RGB-tip ($z' \sim 23.3$) and $z' = 22.1$. In fact, Regions I_g and II_g have similar numbers of bright AGB stars with respect to RGB stars. To demonstrate this point, the ratios of the number of sources in Regions I_g and II_g with z' between 22.1 and 23.3, which are on the AGB, to those with z' between 23.3 and 24.1, which are on the upper RGB, R_{RGB}^{AGB} , were calculated, and the results are given in Table 3. R_{RGB}^{AGB} in Region II_g exceeds that in Region I_g at only slightly

²Age is another parameter that can affect RGB color. However, in §5.3 it is demonstrated that, based on the ratio of bright AGB stars to bright RGB stars, the age distributions of Regions I_g and II_g during intermediate epochs were likely similar, suggesting that the RGBs of Regions I_g and II_g may not have vastly different ages.

more than the $1 - \sigma$ level, indicating that there is not a statistically significant difference in the number of AGB stars with respect to RGB stars. Given the dearth of young main sequence stars in Region II_g, the similar values of R_{RGB}^{AGB} suggests that intermediate age and very young stars in the outer regions of NGC 2403 have different spatial distributions, in the sense that the intermediate age population extends out to larger R_{GC} than the very young bright main sequence stars.

The bolometric LF of AGB stars in Region II_g, constructed from the $(i', r' - i')$ CMD using the procedure discussed in §4, is shown in Figure 15. As in Region I_g, the general shape of the LF is consistent with the Hudon et al. (1989) Field 2 + 3 LF, as well as with the LMC LF from Reid & Mould (1984), providing further evidence that the intermediate age contents of Regions I_g and II_g are similar. The most luminous AGB stars in Region II_g have $M_{bol} = -5.5$, which is fainter than in Region I_g. However, this may simply be a stochastic effect arising from the lower number density of objects in Region II_g.

6. NGC 2403 REGION III_g: THE HALO/OUTER HI DISK

6.1. The LFs, TCDs, and CMDs of NGC 2403 Region III_g

Region III_g samples R_{GC}^{disk} between 15 and 19 kpc, and likely contains stars belonging to the halo and the outer HI disk. The g' and i' LFs of Region III_g, corrected for incompleteness and sources in the control field, are shown in Figure 6. The i' LF of Region III_g differs from those of Regions I_g and II_g in that the onset of the RGB between $i' = 23.5$ and 24.0 is the dominant feature in the Region III_g LF, whereas in Regions I_g and II_g the i' LF more-or-less continuously climbs towards fainter i' . There is a modest population of objects in the i' LF of Region III_g that extend up to 1.5 mag above the RGB-tip, and these are stars evolving on the AGB (§6.3).

Region III_g is dominated by red stars. The $(g' - r', r' - i')$ and $(r' - i', i' - z')$ TCDs of Regions III_g and III_c are compared in Figures 16 and 17. The number density of sources on the $(g' - r', r' - i')$ TCDs of Regions III_g and III_c differ markedly when $g' - r' > 0.5$, while there is an absence of objects with stellar SEDs and $r' - i' < 0$ and $g' - r' < 0$ in Region III_g. As in Regions I_g and II_g, the $(g' - r', r' - i')$ TCD of Region III_g contains stars with $g' - r' < 1.5$ and $r' - i' > 0.4$, which are interpreted as giants that are reddened by dust internal to NGC 2403. The upper envelope of these stars in the Region III_g $(g' - r', r' - i')$ TCD suggests that the maximum internal A_g in Region III_g is comparable to that in Region I_g.

The $(r', g' - r')$, $(i', r' - i')$, and $(z', i' - z')$ CMDs of Regions III_g and III_c are shown

in Figures 18, 19, and 20. There is a modest population of objects in the $(r', g' - r')$ CMD of Region III_g with brightnesses and colors that are appropriate for upper main sequence stars. However, a comparable number of objects is also seen in the same area of the Region III_c $(r', g' - r')$ CMD, indicating that the blue objects in Region III_g likely belong to the foreground or background. There is also an absence of stars in the $(r', g' - r')$ CMD of Region III_g with photometric properties suggestive of Type Ia supergiants. Region III_g is thus free of the massive young stars associated with the classical star-forming disk of NGC 2403.

6.2. RGB Stars in NGC 2403 Region III_g

The $(i', r' - i')$ CMD of Region III_g shows a broad giant branch when $i' > 23.5$, with a blue edge near $r' - i' = -0.1$. The i' LF of stars with $r' - i'$ between 0.0 and 0.7 in Region III_g is shown in Figure 12. A least squares fit to the entries with i' between 23.6 and 24.8 indicates that the power law exponent of the Region III_g LF is 0.60 ± 0.16 , which is similar to that measured in Region I_g, and is not significantly different from that measured in globular clusters.

The $(r' - i')$ color distributions of stars with i' between 24 and 25 in the $(i', r' - i')$ CMDs of Region III_g, corrected for sources in Region III_c, is shown in Figure 13. As in Regions I_g and II_g, the width of the $r' - i'$ distribution in Region III_g is consistent with the observational errors predicted from the artificial star experiments. However, the giant branch in Region III_g has a mean color $\overline{r' - i'} = 0.33 \pm 0.02$, which is significantly smaller than in Regions I_g and II_g. Given that Regions I_g and III_g appear to have comparable amounts of internal differential reddening (§6.1), then the trend of decreasing $\overline{r' - i'}$ with increasing R_{GC} across the GMOS field is likely not due to reddening differences, but instead is a consequence of a radial change in stellar content.

6.3. AGB Stars in NGC 2403 Region III_g

A population of AGB stars is evident above the RGB-tip in the $(i', r' - i')$ and $(z', i' - z')$ CMDs of Region III_g. The AGB sequence on the $(z', i' - z')$ CMD in Figure 20 is morphologically similar to that in Regions I_g and II_g. The z' LF of stars in Region III_g with $i' - z'$ colors between 0 and 1 is shown in Figure 14, and the similarity with the LFs of Regions I_g and II_g is obvious, both in terms of the peak z' brightness, and the shape of the LF. R_{RGB}^{AGB} for Region III_g is listed in Table 3. This quantity is similar to what was

measured in Regions I_g and II_g , indicating that the bright AGB stars are uniformly mixed with fainter stars between $R_{GC}^{disk} = 7$ and 19 kpc in NGC 2403.

The bolometric LF of AGB stars in Region III_g is shown in Figure 15. As was the case for Regions I_g and II_g , there is broad agreement with the Field 2 + 3 LF of Hudon et al. (1989), as well as the Reid & Mould (1984) LMC LF. The most luminous AGB stars in Region III_g have $M_{bol} = -6$, and so the peak luminosity in Region III_g is similar to that in Region I_g . The comparisons in Figures 14 and 15 indicate that Regions I_g , II_g , and III_g experienced similar star-forming histories during the past few Gyr.

7. THE HALO AND OUTER HI DISK OF M33

7.1. The LFs, TCDs, and CMDs of the Outer Regions of M33

With a distance of 0.9 Mpc, M33 is roughly $3 - 4\times$ closer than NGC 2403, and so a single GMOS pointing covers a much smaller chunk of this galaxy than NGC 2403. Therefore, the M33 field samples a range of projected R_{GC} corresponding only to Regions II_g and III_g in NGC 2403. The g' and i' LFs of stars in the M33 field, corrected for incompleteness and sources in the control field, are shown in Figure 6. Contamination from background and foreground sources is substantial at all brightnesses, and the number of objects in the M33 field exceeds that in the control field only when $g' > 22.5$ and $i' > 21$, which corresponds to the upper RGB in M33.

Given that M33 and NGC 2403 have similar morphologies, inclinations, and integrated brightnesses, it might be expected that the number density of red giants in the M33 field should be comparable to that in Regions II_g and III_g . The numbers of stars in the top 1.5 mag in i' of the RGB in the M33 field and the three NGC 2403 regions, based on the LFs in Figure 6, are summarized in Table 4; the density of stars per square arcsec is listed in the last column of this table. Two entries are shown for M33 – the first is the observed density of bright giants, while the second is the density in this field if it were observed at the distance of NGC 2403. The latter quantity falls midway between the densities in NGC 2403 Regions II_g and III_g , confirming that the M33 galaxy field and the two outermost regions of NGC 2403 have similar stellar densities.

Sources in the M33 galaxy and control fields have similar SEDs. This is demonstrated in Figures 16 and 17, where the $(g' - r', r' - i')$ and $(r' - i', i' - z')$ TCDs of sources in these fields are compared. There is a modest excess population of sources in the lower left hand panels of Figures 16 and 17 with respect to the lower right hand panels when $r' - i'$ is between 0 and 1, which is the color interval dominated by M33 RGB stars. Both fields

contain a small number of sources with $g' - r' > 1.5$ and $r' - i' > 0.5$. These objects occur in roughly equal numbers in the M33 galaxy and control fields. A modest population of objects with similar colors are also seen throughout the NGC 2403 control field, in numbers similar to those in the M33 fields. It should be emphasized that the number density of these red objects in the control fields is much lower than that of the highly reddened stars that were identified in NGC 2403 in §4, 5, and 6, and their uniform sky distribution suggests that they are background objects. While this part of the TCD contains high redshift QSOs (e.g. Anderson et al. 2001; Fan 1999), the number density of such sources is much lower than that of the red objects found here.

The $(r', g' - r')$, $(i', r' - i')$ and $(z', i' - z')$ CMDs of the M33 galaxy and control fields are shown in Figures 18, 19, and 20. The high level of contamination from foreground stars and background galaxies is clearly evident when comparing the CMDs of the M33 galaxy and control fields, with only subtle differences between each pair of CMDs. There is a clear absence of bright upper main sequence stars in the $(r', g' - r')$ CMD of M33, and the dominant feature in the CMDs is the RGB. The g' observations of M33 sample stars that are intrinsically fainter than were observed in NGC 2403, and stars on the RGB can be seen in the $(r', g' - r')$ CMD with $r' > 22$ and $g' - r'$ between 0.6 and 1.4.

7.2. RGB Stars in the Outer Regions of M33

The RGB of M33 is evident in the $(i', r' - i')$ CMDs in Figure 19 as a mild increase in stellar density with respect to the control field when $i' > 21$. The LF of stars with $r' - i'$ between 0 and 1 in the M33 field is shown in Figure 12. There is only a modest chance of detecting stars in each 0.2 mag i' bin when $i' < 21$, and the M33 LF in Figure 12 contains gaps near the RGB-tip. The modest number of bright giants in the M33 field prevents us from measuring the power-law exponent of the M33 RGB from these data, so a power-law with the exponent fixed at 0.4, which is the value measured in globular clusters (Davidge et al. 2002), was fit to the LF entries with i' between 21.6 and 22.8, and the result is shown in Figure 12. The fitted relation does a reasonable job of matching points near the RGB-tip at $i' \sim 20.8$, although there appears to be an overall deficiency in the number of giants in the upper 0.5 mag of the RGB in i' .

The small number of bright giants in the M33 galaxy field complicates efforts to investigate the $r' - i'$ color distribution using the procedure employed for NGC 2403; in fact, based on the color distributions in Regions II_g and III_g, it can be anticipated that only a few stars having i' between 21.3 and 22.3 in M33 would be detected per 0.1 mag bin in $r' - i'$ even near the peak of the color distribution. Hence, a wider 0.2 mag binning

was used along the color axis to investigate the $r' - i'$ color distribution in M33, and the results are shown in Figure 13. There is an excess population in the M33 galaxy field with respect to the control field when $r' - i' = 0.5$, and so it is concluded that $\overline{r' - i'} = 0.5 \pm 0.05$ in the M33 galaxy field. RGB stars in the M33 halo field are therefore redder than their counterparts in NGC 2403 Regions II_g and III_g.

7.3. AGB Stars in the Outer Regions of M33

A comparison of the $(z', i' - z')$ CMDs in the lower right and left hand panels of Figure 20 indicates that the M33 galaxy field contains bright AGB stars. The z' LF of objects with $i' - z'$ between 0 and 1 in the M33 field is shown in Figure 14. There are 22 stars in the M33 galaxy field 1 mag above the RGB-tip, whereas 10 sources were detected in this same brightness interval in the control field. Given the modest number of AGB stars, it was decided not to investigate the bolometric LF of these objects. Nevertheless, number counts from the $(i', r' - i')$ CMD indicate that there is a statistically significant number of stars in the M33 GMOS dataset with respect to the control field when $M_{bol} < -4.25$.

Some insight into the relative AGB contents of the outer regions of M33 and NGC 2403 can be obtained from the R_{RGB}^{AGB} statistic, which was computed for the three NGC 2403 fields. The number of AGB stars in M33 were counted with z' between 19.4 and 20.6, while the number of AGB + RGB stars was based on sources having z' between 20.6 and 21.4 to cover the same intrinsic brightness intervals used to find R_{RGB}^{AGB} in NGC 2403. R_{RGB}^{AGB} for M33 is listed in Table 3. The estimated uncertainty in this quantity is substantial due to small number statistics; nevertheless, R_{RGB}^{AGB} in the M33 galaxy field may be slightly larger than in NGC 2403, although the statistical significance is modest. While images covering a much larger field of view will be required to determine the shape of the AGB LF and the luminosity of the AGB-tip in the outer regions of M33, the presence of stars with $M_{bol} < -4$ indicates that, like NGC 2403, the outer regions of M33 contain bright AGB stars.

8. DISCUSSION & SUMMARY

Deep $g'r'i'z'$ images obtained with the GMOS on Gemini North have been used to investigate the brightest RGB and AGB stars in the inner halo and outer disk of the nearby Sc galaxies M33 and NGC 2403. The global properties of these galaxies are similar in many respects, and so it might be expected that the stellar contents of these systems should also be similar. The age distribution of stars in the outer disk of NGC 2403 is discussed in §8.1,

while the metallicities of the M33 and NGC 2403 fields are compared in §8.2. The nature of the bright AGB stars, which have been detected in the outer regions of both galaxies, is discussed in §8.3.

8.1. Comparing the Age Distributions of Stars in the Star-forming Disk of NGC 2403

The stellar content in the outermost regions of disks can be used to probe the physical processes that drive disk evolution. The age distribution of stars is of particular interest, as infall and viscosity may feed star-forming material to the outer disk at late epochs, and spur star formation in regions that previously have not formed stars in large numbers. The age distribution in regions of this nature may then be skewed to more recent values than in the inner disk (e.g. Ferguson & Clarke 2001). Interactions with satellites can also have a major impact on the outer disk, creating disruptions at large radii (e.g. Laurikainen & Salo 2001), and possibly causing the radial expansion of the disk (Aguerre et al. 2001). Tidal stirring may also delay the formation of the outer disk, which has a lower density than the inner disk, and hence is more easily disrupted.

The AGB is a useful evolutionary phase for probing the star-forming history of disks, as it samples stars with ages spanning one Gyr or more. In §4 it was demonstrated that the AGB LF of Region I_g is morphologically similar to that measured by Hudon et al. (1989) in their Fields 2 and 3, and by Reid & Mould (1984) in the LMC. However, to gain insight into the relative age distributions of these regions it is necessary to compare the density of bright AGB stars normalized to other stellar objects that trace the mass distribution. One approach is to scale the AGB LFs according to total stellar density, which can be estimated from surface brightness measurements, such as those obtained by Kent (1987a) for NGC 2403 in the *r*-band; a key assumption is that the M/L ratio does not change markedly from field-to-field.

The *r*-band surface brightness measurements for Region I_g and each of the Hudon et al. (1989) fields, based on the entries in Table II of Kent (1987a), are listed in Table 5. Kent (1987a) did not measure the surface brightness of NGC 2403 along the minor axis beyond 4.8 arcmin from the galaxy center, and so his measurements do not include Region I_g, which is 7.5 arcmin from the center of the galaxy. The surface brightness for Region I_g listed in Table 5 was obtained by extrapolating the minor axis surface brightness measurements between 2.5 and 4.8 arcmin in Table II of Kent (1987a) out to 7.5 arcmin.

The AGB LFs of Region I_g and Hudon et al. (1989) Fields 1 and Fields 2+3 are

compared in Figure 21, where the Hudon et al. (1989) LFs have been scaled to match the surface brightness and areal coverage of Region I_g. The AGB LFs of Region I_g and Fields 2 + 3 agree over the full range of luminosities sampled, indicating identical fractional contributions from luminous AGB stars to the total stellar content. This in turn implies similar age distributions during the time scales probed by AGB stars with $M_{bol} < -4.5$, which corresponds to a few Gyr. Region I_g and Hudon et al. (1989) Fields 2 + 3 sample very different parts of the galaxy, and the agreement between the AGB LFs of such spatially diverse regions indicates that any stochastic influences on the star-forming history in the outer regions of NGC 2403 average out over Gyr timescales.

The surface brightness measurements indicate that the stellar density in Field 1 is $100\times$ that in Region I_g. The AGB LF of Field 1 agrees within the estimated uncertainties with those of Region I_g and Fields 2 + 3 when $M_{bol} < -5$, indicating that luminous AGB stars account for similar fractions of the total r -band light in these areas. Despite this agreement at the bright end, the Field 1 LF falls below those of Region I_g and Fields 2 + 3 when $M_{bol} = -4.5$. This may suggest that the *inner* disk of NGC 2403 experienced a lull in star formation with respect to the *outer* disk within the past few Gyr; in other words, this comparison ostensibly suggests that the age distributions of the inner and outer disks during intermediate epochs differ. However, it is clear from Figure 6 of Hudon et al. (1989) that stars with M_{bol} near -4.5 are at the faint limit of their data, raising the possibility that the absence of stars with $M_{bol} = -4.5$ in Field 1 may be a consequence of incompleteness.

Metallicity gradients are observed in nearby disks (e.g. Zaritsky, Kennicutt, & Huchra 1994; Bell & de Jong 2000), and the gradient in NGC 2403 measured from HII regions is $\Delta[\text{O}/\text{H}]/\Delta r = -0.089 \pm 0.033 \text{ dex kpc}^{-1}$. (Zaritsky, Elston, & Hill 1990). Consequently, at least to the extent that $[\text{O}/\text{H}]$ in HII regions tracks changes in the metallicity of stars formed a Gyr in the past, then $\Delta[\text{O}/\text{H}] = 0.3$ between Region I_g and Hudon et al. (1989) Field 1. Changes in metallicity of this size do not greatly affect the luminosity of the AGB-tip (e.g. Bertelli et al. 1994), suggesting that the difference between the Region I_g and Field 1 LF in Figure 21 at the faint end is likely not a consequence of differences in mean metallicity.

8.2. The Metallicities of the Outer Regions of NGC 2403 and M33

The R_{RGB}^{AGB} entries for NGC 2403 Regions I_g, II_g and III_g in Table 3 are not significantly different, suggesting that these areas have similar age distributions, at least during the past few Gyr. If this is the case, then the region-to-region differences in the mean $r' - i'$ colors of bright RGB stars are most likely due to metallicity. There is as yet no empirical

or theoretical calibration relating $r' - i'$ to metallicity, and so a preliminary calibration is determined here using (1) the cluster giant branch sequences of Da Costa & Armandroff (1990), (2) the relation between $V - I$ and $R - I$ for giants from Bessell (1979), and (3) the transformation between $r' - i'$ and $R - I$ from Fukugita et al. (1996).

In keeping with the $r' - i'$ color distributions shown in Figure 13, the metallicity calibration is based on the $r' - i'$ color one mag in i' below the RGB-tip. The $V - I$ colors of the clusters studied by Da Costa & Armandroff (1990) one mag in I below the RGB-tip are shown in Table 6, along with the $R - I$ and $r' - i'$ colors computed with the Bessell (1979) and Fukugita et al. (1996) relations. Assuming that most of the stars in the NGC 2403 and M33 fields are not subject to significant internal extinction, then the $r' - i'$ and $[\text{Fe}/\text{H}]$ entries in Table 6 indicate that the mean metallicities of NGC 2403 Regions I_g , II_g , and III_g are -0.8 ± 0.1 (random) ± 0.3 (systematic), -1.4 ± 0.1 (random) ± 0.4 (systematic), and -2.2 ± 0.2 (random) ± 0.8 (systematic), respectively. The random errors reflect the statistical uncertainty in $\overline{r' - i'}$, while the systematic errors are due to the ± 0.05 mag uncertainty in the photometric calibration.

The $r' - i'$ colors computed for the calibrating globular clusters in Table 6 rely on color-color relations that are dominated by solar neighborhood stars. This is a potential source of systematic uncertainty, and so the calibration was checked at both the metal-poor and metal-rich ends. The calibration at the metal-poor end was checked directly by comparing the observed $r' - i'$ RGB colors of the metal-poor globular cluster NGC 2419 and the dwarf spheroidal galaxy And V, both of which were discussed by Davidge et al. (2002), with the predicted values. The observed $r' - i'$ RGB colors of these systems one mag in i' below the RGB-tip are listed in the last two rows of Table 6; the uncertainty in the photometric calibration of both systems is ± 0.05 mag (Davidge et al. 2002). And V and NGC 2419 have $[\text{Fe}/\text{H}] = -2.2$, and so should have $r' - i'$ comparable to that of the globular cluster NGC 7078, for which $r' - i' = 0.3$ in Table 6. Given the ± 0.05 mag uncertainty in the photometric calibration of the NGC 2419 and And V data, then the observed RGB colors of these systems are consistent with that of NGC 7078.

The check of the metallicity calibration at the metal-rich end is less direct, and relies on HII regions. Disks contain metallicity gradients and so only HII regions with R_{GC} comparable to Region I_g are considered. There are two HII regions in the Zaritsky et al. (1990) dataset, which are #'s 0 and 29 in their Table I(c), that have R_{GC} close to that of NGC 2403 Region I_g . Source 0 has $\log[\text{OIII}]/\text{H}\beta = 0.91$, whereas for source 29 this ratio is 0.55. Using the calibration of Edmunds & Pagel (1984), these ratios correspond to

$12 + \log[\text{O}/\text{H}] \sim 7.9$ or 8.1 for $\# 0$, and 7.3 or 8.3 for $\# 29$.¹ If $12 + [\text{O}/\text{H}] = 8.9$ in the sun (Anders & Grevesse 1989), then Source $\# 0$ has $[\text{O}/\text{H}] = -0.8$ or -1.0 , whereas for $\# 29$ $[\text{O}/\text{H}] = -0.6$ or -1.6 . The upper $[\text{O}/\text{H}]$ values of these HII regions are in reasonable agreement, whereas this is not the case for the lower values, and so the upper values are adopted here to estimate $[\text{O}/\text{H}]$ in Region I_g . Taking the mean of the upper $[\text{O}/\text{H}]$ values for these two HII regions gives $[\text{O}/\text{H}] \sim -0.7$ for Region I_g , which is 0.3 dex lower than in the LMC (Lequeux et al. 1979; Peimbert & Torres-Peimbert 1974). If the young populations in the LMC have $[\text{M}/\text{H}] = -0.3$ (Harris & Zaritsky 2001) and if the chemical mixtures in the outer disk of NGC 2403 and the LMC are similar, then the young stars in Region I_g would then be 0.3 dex more metal-poor, so that $[\text{M}/\text{H}] = -0.6$. If the chemical evolution of Region I_g has been similar to that of the LMC, which seems reasonable given the similar AGB LFs (§4), then RGB stars in NGC 2403 will be 0.3 dex more metal-poor than the most recently formed stars (Harris & Zaritsky 2001), so that $[\text{M}/\text{H}] = -0.6 + -0.3 = -0.9$ dex, and this is in excellent agreement with what is computed for Region I_g from the $r' - i'$ color of the RGB.

The metallicity measured for the M33 GMOS field from the $r' - i'$ RGB colors is slightly lower than in the corresponding regions of NGC 2403 (i.e. Regions II_g and III_g), but not significantly so. Based on the calibration listed in Table 6 and after accounting for line of sight reddening we find that $[\text{Fe}/\text{H}] = -1.0 \pm 0.3$ (random) ± 0.3 (systematic) for the M33 GMOS field. For comparison, the mean metallicity in NGC 2403 Regions II_g and III_g is -1.8 ± 0.6 ; hence, there is not a statistically significant difference between the two galaxies.

The metallicity measured from our M33 galaxy field is larger at the 1.4σ level than that measured by MK86 in a field with a projected distance of 7 kpc from the center of M33. MK86 found that stars in this field have a mean metallicity comparable to that of the $[\text{Fe}/\text{H}] \sim -2.2$ cluster M92, and assigned an error of ± 0.8 dex due to uncertainties in the photometric zeropoint. A metallicity as low as that measured by MK86 for the halo of M33 (and that measured here for NGC 2403 Region III_g) is not consistent with what might be expected from the globular cluster system of M33. CS88 studied the ages and metallicities of M33 clusters, and the six oldest clusters in their compilation, which are those with $\log(t) \geq 9.6$ in their Table VIII, have $[\text{Fe}/\text{H}]$ between -1.2 and -2.2 , with a mean $[\text{Fe}/\text{H}] = -1.55 \pm 0.15$. Chandar et al. (2002) also find that the majority of halo globular clusters in M33 have $[\text{Fe}/\text{H}]$ between -1 and -2 . Given that globular clusters tend to be more metal-poor than their surroundings (e.g. Forbes et al. 1996; Da Costa & Mould 1988), then

¹The relation between $\log[\text{OIII}]/\text{H}\beta$ and $\log(\text{O}/\text{H})$ is multi-valued (Edmunds & Pagel 1984), and so two $[\text{O}/\text{H}]$ values are quoted here.

it can be anticipated that the halo of M33 should contain a significant field population with $[\text{Fe}/\text{H}] \geq -1.6$, which is contrary to the MK86 metallicity.

Cuillandre, Lequeux, & Loinard (1999) investigated the stellar content of a large field in the south east portion of M33, which included the area studied by MK86. Based on the RGB sequence in the $(I, V - I)$ CMD, Cuillandre et al. (1999) conclude that $[\text{Fe}/\text{H}] = -1$ with ± 0.2 dex error due to uncertainties in the distance to M33. The width of the RGB is consistent with a $\pm 0.4 - 0.5$ dex spread in metallicity, although the field may be contaminated by outer disk stars with a range of ages. Nevertheless, the mean metallicity measured by Cuillandre et al. (1999) is consistent with that found in the GMOS M33 field and in Region I_g of NGC 2403.

8.3. Intermediate-age Populations Outside of the Present-Day Star-Forming Disks of NGC 2403 and M33

Observations of edge-on spiral galaxies indicate that stellar disks terminate on spatial scales of roughly 1 kpc (van der Kruit & Searle 1982; de Grijs, Kregel, & Wesson 2001 and references therein), which corresponds to ~ 1 arcmin at the distance of NGC 2403. The radius at which disk truncation occurs is still a matter of some debate, and factors such as environment likely play a significant role. van der Kruit & Searle (1982) find that the disk cut-off occurs typically at $(4.2 \pm 0.5) \times h_r$, where h_r is the radial scale length of the disk, whereas Pohlen, Dettmar, & Lütticke (2000) find a mean cut-off radius of $(2.9 \pm 0.7) \times h_r$. The disks in NGC 2403 and M33 have scale lengths of 2.1 – 2.2 kpc (Kent 1987a,b), and so the stellar disks of these systems might be expected to truncate at $R_{GC}^{disk} \sim 6 - 9$ kpc. This is roughly consistent with the outer boundary of Region I_g , where there is a clear drop in the blue main sequence component.

The presence of bright AGB stars in NGC 2403 Region III_g indicates that star formation occurred over a much larger range of R_{GC}^{disk} in NGC 2403 during intermediate epochs than in the present-day star-forming disk. Luminous AGB stars are also present at large R_{GC} in the M33 GMOS data (§7), as well as in the CMD presented in Figure 4 of MK86, which contains stars that are redder and brighter than the RGB-tip. In fact, there are 29 sources in the MK86 dataset in the I interval 1 mag above the RGB-tip, and number counts in the M33 GMOS control field predict that only 6 of these are background or foreground sources.

The detection of bright AGB stars at large R_{GC} in NGC 2403 and M33 is perhaps not surprising, as it has been known for some time that the globular cluster systems of

these galaxies contain intermediate-age objects at large R_{GC} . Battistini et al. (1984) used $B - V$ colors to identify three classes of clusters in NGC 2403. While the majority of blue clusters have small R_{GC} , one (C3) is 7 arcmin ($R_{GC}^{LOS} \sim 6.4$ kpc) from the galaxy center, while another (F2) is 15 arcmin ($R_{GC}^{LOS} \sim 14$ kpc) from the center. The M33 globular cluster system also contains young and intermediate age clusters (CS88, Sarajedini et al. 1998, Chandar et al. 2002, Ma et al. 2002a,b), and many of these are located at large R_{GC} (e.g. Figure 3 of CS88), placing them outside of the main body of the disk. The presence of intermediate age globular clusters with halo-like kinematics in M33 (e.g. Chandar et al. 2002) indicates that the giant molecular clouds that are thought to be the birth sites of these systems (e.g. Harris & Pudritz 1994; Ashman & Zepf 2001) were present at large R_{GC} in this galaxy or its satellites during intermediate epochs.

M33 and NGC 2403 are not unique among nearby spiral galaxies in having intermediate-age populations outside of their active star-forming disks, as Ferguson & Johnson (2001) found main sequence stars with a mass of $3 M_{\odot}$ in a semi-major axis field 30 kpc from the center of M31. However, the outer regions of M31 do not contain the young and intermediate age globular clusters seen in M33 and NGC 2403. This could indicate that the intermediate-age field populations in the outer regions of M31 have a smaller fractional size than in M33 and NGC 2403, or that there is a difference in origin.

Did the luminous AGB stars in NGC 2403 and M33 form in situ, or did they form in a satellite (or satellites) that was (were) accreted during recent epochs? An accretion origin is attractive because satellite systems at moderately large R_{GC} may be able to retain gas up to recent epochs, and hence provide a reservoir of material for star formation. However, late satellite accretion does not provide an ironclad explanation for the origin of bright AGB stars in the outer regions of NGC 2403 and M33. In particular, if the halo potentials of these galaxies are near spherical, as predicted by simulations (e.g. Bekki & Chiba 2001), then the rate of precession will be modest and tidal streams will not smear markedly with time (Ibata et al. 2001). The AGB stars in M33 and NGC 2403 appear not to be in stream-like structures. In fact, the uniformity of the quantity R_{RGB}^{AGB} , defined in §5, with radius in NGC 2403, even when including fields that sample the young disk, requires that the tidally accreted material would have to be well-mixed within the host galaxy.

The tendency for R_{RGB}^{AGB} not to change with radius in NGC 2403 suggests that the bright AGB stars formed in situ. Are the conditions in the outer regions of NGC 2403 and M33 at the present epoch such that star formation can occur? Star formation is suppressed when the ratio of the local gas density to the density required to maintain stability based on the Toomre criterion drops below ~ 0.7 (Kennicutt 1989; Martin & Kennicutt 2001). Figure 8 of Martin & Kennicutt (2001) indicates that the gas density in NGC 2403 exceeds

that required for star formation out to large radii, so it presumably should not be difficult to spur star formation at large R_{GC} in NGC 2403.

Figure 13 of Martin & Kennicutt (2001) indicates that M33 does not contain an extended gas envelope with a density that is high enough to form stars according to the standard Kennicutt (1989) threshold. Nevertheless, the HI envelope of M33 contains structure (Corbelli, Schneider, & Salpeter 1989), and star formation can occur in regions where the mean gas density is lower than the critical value (e.g. Ferguson et al. 1998), since random perturbations and spiral density waves can cause localized gas compression, and thereby trigger star formation (e.g. Kennicutt 1989, Ferguson et al. 1998, Martin & Kennicutt 2001). If star formation is triggered by such processes then it might be anticipated that the star-forming history in the outermost regions of a galaxy will be less continuous than at smaller radii. This being said, the similar AGB contents of Regions I_g, II_g, and III_g and Hudon et al. (1989) Fields 2 + 3 indicates that star formation in the outermost region of NGC 2403 has been coherent over galaxy-wide spatial scales when averaged over Gyr time scales.

Both Regions II_g and III_g sample the outer HI disk of NGC 2403, and if the intermediate age stars formed from material from this gas reservoir then they might be expected to fall along the plane of the HI disk and have disk-like kinematics. Investigations of the number density of bright AGB stars in different fields in the outer regions of M33 and NGC 2403, coupled with spectroscopic studies to obtain kinematic information, will ultimately provide the information needed to chart the spatial distribution of these objects. However, there are tantalizing indications that the intermediate age globular clusters in late-type spiral galaxies do not belong to a disk. For example, Chandar et al. (2002) find that many of the intermediate age clusters in M33 do not have disk-like kinematics, but appear to belong to a pressure-supported halo. Moreover, Kissler-Patig et al. (1999) have found a population of blue clusters well off of the disk of the edge-on Sc galaxy NGC 5907.

Bekki & Chiba (2001) investigated the formation of halos in the context of CMD-based models, and suggested that the amount of halo flattening depends on host galaxy morphology, in the sense that late-type systems have flatter inner halos. These simulations predict that a flattened inner halo in late-type spirals (1) forms well after the more spherically distributed outer halo, and (2) contains younger stars, which formed during the dissipative merging of gas-rich satellites, than the outer halo. The luminous AGB stars that occur outside of the actively star-forming disks of M33 and NGC 2403 may be the remnants of the final collapse phase of these systems. Such a delayed formation model is consistent with the morphological evolution of Sc galaxies with redshift, in that these systems are observed to become increasingly chaotic at intermediate redshifts (van den Bergh 2002).

8.4. Future Work

There is an obvious need to investigate additional fields sampling a larger range of R_{GC} in M33 and NGC 2403, as studies of the spatial distribution of AGB stars in the outer regions of these galaxies will provide clues about the origin of intermediate age populations at large R_{GC} . Photometric and spectroscopic surveys of the M33 and NGC 2403 globular cluster systems will also provide additional insight into the star-forming histories of the outer regions of these galaxies. Contamination from background galaxies and foreground stars, which our data demonstrates is already significant at the brightness of the RGB-tip when $R_{GC}^{LOS} = 9$ kpc in M33, will become an even greater issue at larger radii, and spectra will become increasingly important for identifying stars that belong to either galaxy. While spectroscopic observations of RGB-tip stars in Local Group galaxies are feasible with existing 8 – 10 metre telescopes, studies of more distant systems, like NGC 2403, must await 20 – 30 metre facilities. Nevertheless, infrared colors and narrow-band photometry, which can be obtained for these stars using 8 – 10 metre telescopes, offer one means of distinguishing between halo and contaminating stars in an efficient manner; not only are stars near the AGB-tip relatively bright in JHK , but the near-infrared SEDs of dwarfs and giants differ, so that foreground stars can be identified.

The late-type Sculptor group galaxies NGC 55 and NGC 247 are potentially interesting targets for surveys of bright AGB stars at large R_{GC} . Like M33 and NGC 2403, NGC 55 and NGC 247 are Sc galaxies. NGC 55 and NGC 247 have distances that are intermediate between M33 and NGC 2403, and are located near the South Galactic Pole, so that there will be lower contamination from foreground stars than in either NGC 2403 or M33. NGC 55 is also viewed almost edge-on, thereby simplifying the task of studying the spatial distribution of stars away from the plane of the disk.

It is a pleasure to thank Sidney van den Bergh, Harvey Richer, and Jean-Rene Roy for commenting on an earlier version of the manuscript. Sidney van den Bergh also suggested that the globular cluster systems of M33 and NGC 2403 be considered when discussing the stellar contents of these galaxies. Thanks are also extended to Inger Jørgensen, who co-ordinated the GMOS SV process, and to an anonymous referee, who made a number of comments that improved the paper.

Galaxy	μ_0	E(B–V) ^a
NGC 2403	27.51 ± 0.24 ^b	0.046 (0.052)
M33	$24.81^{+0.15}_{-0.11}$ ^c	0.057 (0.079)

Table 1: Adopted Distances and Reddenings

^aFrom Schlegel, Finkbeiner, & Davis (1998). The entry in brackets is the reddening of the control field for that galaxy.

^bFreedman & Madore (1988)

^cKim et al. (2002)

Field	RA (J2000)	Dec (J2000)	Filter	Exposure Time (seconds)	FWHM (arcsec)
M33	01:36:27.6	+30:16:41	g'	6×60	1.16
			r'	6×60	0.94
			i'	6×60	0.87
			z'	6×60	0.65
M33 control	01:45:01.2	+27:32:45	g'	6×60	0.94
			r'	6×60	0.94
			i'	6×60	0.87
			z'	6×60	0.65
NGC 2403	07:37:51.1	+65:43:06	g'	6×900	1.23
			r'	6×400	0.87
			i'	6×400	0.80
			z'	12×200	0.72
NGC 2403 control	07:52:32.6	+61:23:32	g'	2×900	1.02
			r'	6×400	0.65
			i'	6×400	0.65
			z'	12×200	0.87

Table 2: Details of the Observations

Region	R_{RGB}^{AGB}
I _g	0.22 ± 0.07
II _g	0.44 ± 0.18
III _g	0.23 ± 0.12
M33	0.9 ± 0.5

Table 3: The ratio of bright AGB stars to upper RGB stars in each field

Field	Number of stars	Density (arcsec ⁻²)
Region I _g	388.5 ± 30.9	$(6.5 \pm 0.5) \times 10^{-6}$
Region II _g	187.5 ± 36.3	$(2.5 \pm 0.5) \times 10^{-6}$
Region III _g	166.0 ± 39.4	$(1.6 \pm 0.4) \times 10^{-6}$
M33	31.0 ± 12.7	$(1.3 \pm 0.5) \times 10^{-7}$
M33 _{NGC2403}	–	$(2.1 \pm 0.8) \times 10^{-6}$

Table 4: The density of stars within 1.5 mag of the RGB-tip

Field	Surface Brightness (r -band) ^a
Region I _g	26.6
Field 1	21.8
Fields 2 + 3	24.3

Table 5: r -band surface brightness measurements for Region I_g and the Hudon et al. fields

^aFrom Kent (1987)

Object	[Fe/H]	$(V - I)_{RGB}$	$(R - I)_{RGB}$	$(r' - i')_{RGB}$
N104	−0.7	1.54	0.74	0.50
N1851	−1.3	1.38	0.65	0.41
N6752	−1.5	1.28	0.60	0.36
N7089	−1.6	1.26	0.59	0.35
N7078	−2.2	1.17	0.54	0.30
And V	−2.2	—	—	0.36
N2419	−2.2	—	—	0.34

Table 6: $r' - i'$ colors one mag below the RGB-tip for selected globular clusters and the dwarf spheroidal galaxy And V

REFERENCES

- Aguerri, J. A. L., Balcells, M., & Peletier, R. F. 2001, *A&A*, 367, 428
- Anders, E., & Grevesse, N. 1989, *Ge & Cosm Acta* 53, 197
- Anderson, S. F. et al. 2001, *AJ*, 122, 503
- Ashman, K. M., & Zepf, S. E. 2001, *AJ*, 122, 1888
- Battistini, P., Bònoli, F., Federici, L., Fusi Pecci, F., & Kron, R. G. 1984, *A&A*, 130, 162
- Bekki, K., & Chiba, M. 2001, *ApJ*, 558, 666
- Bell, E. F., & de Jong, R. S. 2000, *MNRAS*, 312, 497
- Bessell, M. S. 1979, *PASP*, 91, 589
- Bessell, M. S., & Wood, P. R. 1984, *PASP*, 96, 247
- Bica, E., Barbuy, B., Ortolani, S. 1991, *ApJ*, 382, L15
- Binney, J., Dehnen, W., & Bertelli, G. 2000, *MNRAS*, 318, 658
- Bullock, J. S., Kravtsov, A. V., & Weinberg, D. H. 2001, *ApJ*, 548, 33
- Buonanno, R., Corsi, C. E., Pulone, L., Fusi Pecci, F., & Bellazzini, M. 1998, *A&A*, 333, 505
- Burstein, D., Faber, S. M., Gaskell, C. M., & Krumm, N. 1984, *ApJ*, 287, 586
- Caldwell, N., Armandroff, T. E., Da Costa, G. S., & Seitzer, P. 1998, *AJ*, 115, 535
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Chandar, R., Bianchi, L., Ford, H. C., & Sarajedini, A. 2002, *ApJ*, 564, 712
- Chiba, M., & Beers, T. C. 2001, *ApJ*, 549, 325
- Christian, C. A., & Schommer, R. A. 1988, *AJ*, 95, 704 (CS88)
- Cole, A. A., Smecker-Hane, T. A., & Gallagher, J. S. 2000, *AJ*, 120, 1808
- Corbelli, E., Schneider, S. E., & Salpeter, E. E. 1989, *AJ*, 97, 390
- Côté, P., Marzke, R. O., West, M. J., & Minniti, D. 2000, *ApJ*, 533, 869
- Crampton, D. et al. 2000, *SPIE*, 4008, 114
- Cuillandre, J-C, Lequeux, J., & Loinard, L. 1999 in *IAU Symposium 192 ‘The Stellar Content of Local Group Galaxies’*, P. Whitelock & R. Cannon ed.
- Da Costa, G. S., & Armandroff, T. E. 1990, *AJ*, 100, 162
- Da Costa, G. S., & Mould, J. R. 1988, *ApJ*, 334, 159

- Davidge, T. J. 1990, *ApJ*, 351, L37
- Davidge, T. J. 2002, *PASP*, 114, 1206
- Davidge, T. J., & Courteau, S. 2002, *AJ*, 123, 1438
- Davidge, T. J., Da Costa, G. S., Jørgensen, I., & Allington-Smith, J. R. 2002, *AJ*, 124, 886
- de Grijs, R., Kregel, M., & Wesson, K. H. 2001, *MNRAS*, 324, 1074
- Durrell, P. R., Harris, W. E., & Pritchett, C. J. 1994, *AJ*, 108, 2114
- Durrell, P. R., Harris, W. E., & Pritchett, C. J. 2001, *AJ*, 121, 2557
- Edmunds, M. G., & Pagel, B. E. J. 1984, *MNRAS*, 211, 507
- Fan, X. 1999, *AJ*, 117, 2528
- Ferguson, A. M. N., & Clarke, C. J. 2001, *MNRAS*, 325, 781
- Ferguson, A. M. N., & Johnson, R. A. 2001, *ApJ*, 559, L13
- Ferguson, A. M. N., Wyse, R. F. G., Gallagher, J. S., & Hunter, D. A. 1998, *ApJ*, 506, L19
- Forbes, D. A., Franx, M., Illingworth, G. D., & Carollo, C. M. 1996, *ApJ*, 467, 126
- Fraternali, F., van Moorsel, G., Sancisi, R., & Oosterloo, T. 2002, *AJ*, 123, 3124
- Freedman, W. L., & Madore, B. F. 1988, *ApJ*, 332, L63
- Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Simasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
- Garnett, D. R., Shields, G. A., Skillman, E. D., Sagan, S. P., & Dufour, R. J. 1997, *ApJ*, 489, 63
- Grillmair, C. J. et al. 1996, *AJ*, 112, 1975
- Harris, G. L. H., & Harris, W. E. 2000, *AJ*, 120, 2423
- Harris, W. E., & Harris, G. L. H. 2001, *AJ*, 122, 3065
- Harris, J., & Zaritsky, D. 2001, *ApJS*, 136, 25
- Harris, W. E., & Pudritz, R. E. 1994, *ApJ*, 429, 177
- Helmi, A., & White, S. D. M. 1999, *MNRAS*, 307, 495
- Hudon, J. D., Richer, H. B., Pritchett, C. J., Crabtree, D., Christian, C. A., & Jones, J. 1989, *AJ*, 98, 1265
- Humphreys, R. M., & McElroy, D. B. 1984, *ApJ*, 284, 565
- Ibata, R., Lewis, G. F., Irwin, M., Totten, E., & Quinn, T. 2001a, *ApJ*, 551, 294
- Ibata, R., Irwin, M., Lewis, G., Ferguson, A. M. N., & Tanvir, N. 2001b, *Nature*, 412, 49

- Karachentsev, I. D. et al. 2000, *A&A*, 363, 117
- Kennicutt, R. C. 1989, *ApJ*, 344, 685
- Kent, S. M. 1987a, *AJ*, 93, 816
- Kent, S. M. 1987b, *AJ*, 94, 306
- Kim, M., Kim, E., Lee, M. G., Sarajedini, A., & Geisler, D. 2002, *AJ*, 123, 244
- Kissler-Patig, M., Ashman, K. M., Zepf, S. E., & Freeman, K. C. 1999, *AJ*, 118, 197
- Knox, R. A., Hawkins, M. R. S., & Hambly, N. C. 1999, *MNRAS*, 306, 736
- Krisciunas, K., Margon, B., & Szkody, P. 1998, *PASP*, 110, 1342
- Laurikainen, E., & Salo, H. 2001, *MNRAS*, 324, 685
- Lee, M. G., Kim, M., Sarajedini, A., Geisler, D., & Gieren, W. 2002, *ApJ*, 565, 959
- Leggett, S. K., Ruiz, M. T., & Bergeron, P. 1998, *ApJ*, 497, 294
- Lenz, D. D., Newberg, H. J., Rosner, R., Richards, G. T. & Stoughton, C. 1998, *ApJS*, 119, 121
- Lequeux, J., Peimbert, M., Rayo, J. F., Serrano, A., & Torres-Peimbert, S. 1979, *A&A*, 80, 155
- Liu, W. M., & Chaboyer, B. 2000, *ApJ*, 544, 818
- Ma, J., Zhou, X., Chen, J., Wu, H., Jiang, Z., Xue, S., & Zhu, J. 2002a, *A&A*, 385, 404
- Ma, J. Xu, Z., Chen, J., Wu, H., Jiang, Z., Xue, S., & Zhu, J. 2002b, *AJ*, 123, 3141
- Majewski, S. R., Munn, J. A., & Hawley, S. L. 1994, *ApJ*, 427, L37
- Martin, C. L., & Kennicutt, R. C. 2001, *ApJ*, 555, 301
- Martinez-Delgado, D., Aparicio, A., Angeles Gomez-Flechoso, M., & Carrera, R. 2001, *ApJ*, 549, L199
- McCall, M. L., Rybski, P. M., & Shields, G. A. 1985, *ApJS*, 57, 1
- Mould, J., & Kristian, J. 1986, *ApJ*, 305, 591 (MK86)
- O'Donnell, J. E. 1994, *ApJ*, 422, 158
- Oswalt, T. D., Smith, J. A., Wood, M. A., & Hintzen, P. 1996, *Nature*, 382, 692
- Peimbert, M., & Torres-Peimbert, S. 1974, *ApJ*, 193, 327
- Petersen, L., & Gammelgaard, P. 1996, *A&A*, 308, 49
- Pierce, M. J., & Tully, R. B. 1992, *ApJ*, 387, 47
- Pohlen, M., Dettmar, R.-J., & Lütticke, R. 2000, *A&A*, 357, L1

- Reid, N., & Mould, J. 1984, ApJ, 284, 98
- Rosenberg, A., Saviane, I., Piotto, G., & Aparicio, A. 1999, AJ, 118, 2306
- Saito, Y., & Iye, M. 2000, ApJ, 535, L95
- Sarajedini, A., Chaboyer, B., & Demarque, P. 1997, PASP, 109, 1321
- Sarajedini, A., Geisler, D., Harding, P., & Schommer, R. 1998, ApJ, 508, L37
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, ApJ, 500, 525
- Schneider, D. P., et al. 2001, AJ, 121, 1232
- Stetson, P. B. 1987, PASP, 99, 191
- Stetson, P. B., & Harris, W. E. 1988, AJ, 96, 909
- Tully, R. B., & Fouqué, P. 1985, ApJS, 58, 67
- van den Bergh, S. 2002, PASP, 114, 797
- VandenBerg, D. A. 1992, ApJ, 391, 685
- van der Kruit, P. C., & Searle, L. 1981, A&A, 95, 105
- van der Kruit, P. C., & Searle, L. 1982, A&A, 110, 61
- Weil, M. L., Eke, V. R., & Efstathiou, G. 1998, MNRAS, 300, 773
- Zaritsky, D. 1999, AJ, 118, 2824
- Zaritsky, D., Elston, R., & Hill, J. M. 1990, AJ, 99, 1108
- Zaritsky, D., Kennicutt, R. C., & Huchra, J. P. 1994, ApJ, 420, 87
- Zurek, W. H., Quinn, P. J., & Salmon, J. K. 1988, ApJ, 330, 519

FIGURE CAPTIONS

Fig. 1.— A 20×20 arcmin section of the Digital Sky Survey (DSS) centered on NGC 2403. The box shows the location of the 5.5×5.5 arcmin GMOS field. Although not obvious from this figure, the lower right hand corner of this field contains bright young main sequence stars belonging to the disk of NGC 2403. North is at the top, and East is to the left.

Fig. 2.— A 40×40 arcmin section of the DSS near M33. The box shows the location of the 5.5×5.5 arcmin GMOS field. North is at the top, and East is to the left. The area observed by MK86 is near the center of the figure.

Fig. 3.— The i' images of the NGC 2403 galaxy (top panel) and control (bottom panel) fields. Each image covers approximately 5.5×5.5 arcmin, with North at the top and East to the left. The three regions used to analyze the stellar content of this field are indicated. The dwarf irregular galaxy Kar 50, which was discussed by Davidge (2002), is labelled in the lower panel.

Fig. 4.— The i' images of the M33 galaxy (top panel) and control (bottom panel) fields. Each image covers approximately 5.5×5.5 arcmin, with North at the top and East to the left.

Fig. 5.— The completeness fractions C in g' and i' , as computed from artificial star experiments. C is the ratio of detected artificial stars to the total number inserted into each data frame per 0.5 mag interval. The completeness curves for the outer regions of NGC 2403 and M33 are shown as solid lines, while the corresponding curves for the control fields are shown as dashed lines. The NGC 2403 data go deeper than the M33 data because of the difference in total exposure time (Table 1).

Fig. 6.— The g' and i' LF of sources in the outer regions of NGC 2403 and M33. $N_{0.5}$ is the difference between the number of sources per 0.5 mag interval in the galaxy and control fields, corrected for incompleteness; the dotted line indicates $N_{0.5} = 0$. The g' LFs show sources detected in both g' and r' , while the i' LFs show sources detected in both r' and i' . The error bars reflect uncertainties due to Poisson statistics and the completeness corrections.

Fig. 7.— The $(g' - r', r' - i')$ TCDs of NGC 2403 Regions I and II in the galaxy and control fields. The dotted and dashed lines show the loci of the $\log(g)=2.5$ and 4.5 solar metallicity models of Lenz et al. (1998) while the solid line is the stellar sequence from Figure 3 of Schneider et al. (2001). The reddening vector, which has a length corresponding to $A_{g'} = 1$ mag, follows the relations listed in Table 6 of Schlegel et al. (1998), which in turn are taken from the Cardelli, Clayton, & Mathis (1989) and O'Donnell (1994) reddening curves. Note

that there are a number of sources in Regions I_g and II_g with $r' - i' > 0.4$ and $g' - r' > 1.5$, and objects with corresponding colors are largely absent in Regions I_c and II_c . These red objects are interpreted as giants that are subject to reddening internal to NGC 2403.

Fig. 8.— The $(r' - i', i' - z')$ TCDs of NGC 2403 Regions I and II in the galaxy and control fields. The dotted and dashed lines show the loci of the $\log(g)=2.5$ and 4.5 solar metallicity models of Lenz et al. (1998) while the solid line is the stellar sequence from Figure 3 of Schneider et al. (2001). The reddening vector, which has a length corresponding to $A_{g'} = 1$ mag, follows the relations listed in Table 6 of Schlegel et al. (1998), which in turn are taken from the Cardelli et al. (1989) and O'Donnell (1994) reddening curves.

Fig. 9.— The $(r', g' - r')$ CMDs of NGC 2403 Regions I and II in the galaxy and control fields. The dashed lines in the left hand panel are the main sequence and supergiant Ia sequence defined by stars in the Galaxy and the LMC, shifted to match the distance of NGC 2403 with a correction for foreground reddening only. The solid lines are the same sequences, but with an assumed internal extinction of $A_{g'} = 1$ mag for NGC 2403.

Fig. 10.— The $(i', r' - i')$ CMDs of NGC 2403 Regions I and II in the galaxy and control fields. Both Regions I_g and II_g contain an excess population of sources with respect to the corresponding control fields when $i' > 23.5$, due to stars evolving on the RGB in NGC 2403.

Fig. 11.— The $(z', i' - z')$ CMDs of NGC 2403 Regions I and II in the galaxy and control fields. The dashed line shows $i' = 23.5$, which corresponds to the approximate brightness of the RGB-tip with no internal extinction assumed for NGC 2403, while the solid line shows $i' = 24$, which corresponds to the RGB-tip with an assumed internal extinction $A_{g'} = 1$ mag. Regions I_g and II_g contain an excess population of sources with respect to the corresponding control fields when $z' > 21.5$, and the majority of these are AGB stars in NGC 2403. Aside from a difference in the number of stars, the overall morphologies of the $(z', i' - z')$ CMDs of Region I_g and II_g are not greatly different, suggesting similar AGB contents.

Fig. 12.— The i' LFs of RGB stars in the outer regions of NGC 2403 and M33. $N_{0.2}$ is the difference between the number of stars per 0.2 mag i' interval in the galaxy and control fields in these color intervals, corrected for incompleteness. The errorbars reflect the uncertainties due to Poisson statistics and the completeness corrections. The dotted lines in the top three panels show power laws that have been fit to the data in each NGC 2403 Region when $i' > 23.6$, while the dotted line in the bottom panel shows a power-law with an exponent of 0.4 that has been fit to the M33 LF when i' is between 21.6 and 22.8.

Fig. 13.— The $r' - i'$ distributions of sources on the upper RGB in NGC 2403 and M33. The $r' - i'$ distributions for NGC 2403 were computed using stars with i' between 24 and

25, while the distribution for M33 uses stars with i' between 21.3 and 22.3. The solid lines show the difference between the color distributions in the galaxy and control fields in these brightness intervals, while the error bars show the uncertainties due to Poisson statistics. $N_{0.1}$ and $N_{0.2}$ are the numbers of stars per 0.1 mag and 0.2 mag intervals in $r' - i'$. The dotted lines in the top three panels are gaussians with $\sigma = \pm 0.197$ mag, which is the photometric error predicted by the artificial star experiments, centered on $\overline{r' - i'}$ and scaled to match the number of sources with $r' - i'$ between 0.3 and 0.7 in each region in NGC 2403. Note the excellent agreement with the observed distributions. The dotted line in the bottom panel is a gaussian with $\sigma = \pm 0.03$ mag, which is the photometric error predicted by the artificial star experiments for the M33 data, and $\overline{r' - i'} = 0.5$. The gaussian in the bottom panel has been scaled to match the number of sources in the $r' - i' = 0.5$ bin.

Fig. 14.— The z' LFs of stars with $i' - z'$ between 0 and 1 in the outer regions of NGC 2403 and M33. $N_{0.2}$ is the difference between the number of stars per 0.2 mag z' interval in the galaxy and control fields, corrected for incompleteness. The errorbars reflect the uncertainties due to Poisson statistics and the completeness corrections. The approximate brightness of the RGB-tip is indicated with a vertical line in each panel; sources fainter than the RGB-tip are a combination of RGB and AGB stars, while sources brighter than the RGB-tip are on the AGB. Note that there is a significant number of bright AGB stars in each field.

Fig. 15.— The bolometric LFs of AGB stars in the outer regions of NGC 2403. $N_{0.5}$ is the difference between the number of stars per 0.5 mag M_{bol} interval in the galaxy and control fields, corrected for incompleteness. The errorbars include the uncertainties due to Poisson statistics and the completeness corrections. The LFs were constructed by first transforming i' and $r' - i'$ into I_{KC} and $(R - I)_{KC}$ using the procedure described in the text, and then applying the criteria for identifying AGB stars discussed by Hudon et al. (1989). Bolometric corrections were computed from Equation 2 of Bessell & Wood (1984). The dotted line in each panel is the AGB LF of Hudon et al. (1989) NGC 2403 Fields 2 and 3, while the dashed-dotted line is the LF of AGB stars in the LMC from Reid & Mould (1984). The Hudon et al. and Reid & Mould LFs have been scaled to match the number of objects in each region of NGC 2403 with M_{bol} between -6.2 and -4.5 , while the Hudon et al. (1989) LF has also been shifted by 0.1 mag to match the distance modulus adopted in the current study. Note that both the Hudon et al. and Reid & Mould LFs match the current observations within the estimated uncertainties. The approximate magnitude of the RGB-tip is indicated with the vertical line in each panel; sources fainter than the RGB-tip are a combination of RGB and AGB stars, while sources brighter than the RGB-tip are on the AGB.

Fig. 16.— The $(g' - r', r' - i')$ TCDs of the galaxy and control fields for NGC 2403 Region

III and M33. See Figure 7 for additional details.

Fig. 17.— The $(r' - i', i' - z')$ TCDs of the galaxy and control fields for NGC 2403 Region III and M33. See Figure 8 for additional details.

Fig. 18.— The $(r', g' - r')$ CMDs of the galaxy and control fields for NGC 2403 Region III and M33. The dashed lines in the upper panels are the main sequence and supergiant Ia sequence defined by stars in the Galaxy and the LMC, shifted to match the distance of NGC 2403 for two reddenings (foreground only, and foreground with an internal extinction $A_{g'} = 1$ mag). The dashed lines in the lower panel are the main sequence and supergiant Ia sequences shifted to the distance of M33, with only foreground reddening. See Figure 9 for additional details.

Fig. 19.— The $(i', r' - i')$ CMDs of the galaxy and control fields for NGC 2403 Region III and M33. The majority of stars with $i' > 23.5$ in Region III_g are evolving on the RGB, while those with $i' < 23.5$ are evolving on the AGB. The M33 halo field contains a modest excess population of sources with respect to the control field when $i' > 21$, which is due to the RGB.

Fig. 20.— The $(z', i' - z')$ CMDs of the galaxy and control fields for NGC 2403 Region III and M33. The dashed line in the upper panels marks $i' = 23.5$, which is the approximate brightness of the RGB-tip in NGC 2403 with only foreground extinction, while the solid line marks $i' = 24$, which is the RGB-tip with an assumed internal extinction of $A_{g'} = 1$ mag. The overall morphologies of the $(z', i' - z')$ CMDs of NGC 2403 Regions I_g, II_g, and III_g are very similar. The dashed line in the lower panels marks $i' = 20.8$, which is the approximate brightness of the RGB-tip in M33. The M33 halo field contains an excess population of sources with respect to the control field when $z' < 20$, and the majority of these are RGB stars. The M33 field also contains a modest excess population of sources above the RGB-tip, which are stars evolving on the AGB.

Fig. 21.— The LF of AGB stars in Region I_g (solid line), corrected for foreground and background contamination, compared with the AGB LFs of Hudon et al. (1987) Fields 1 (dashed line) and 2 + 3 (dotted line). The approximate magnitude of the RGB-tip is indicated with the vertical line; sources fainter than the RGB-tip are a combination of RGB and AGB stars, while sources brighter than the RGB-tip are on the AGB. The Hudon et al. (1989) LFs have been shifted along the vertical axis to match the surface brightness in Region I_g using the measurements listed in Table 5 and also to correct for differences in areal coverage. The Hudon et al. LFs have also been shifted by 0.1 mag along the horizontal axis to account for the different distance modulus adopted for the current work. Note that the 3 LFs agree within the estimated errors when $M_{bol} < -5$, indicating that the space density

of luminous AGB stars, when normalised to the total stellar content in each field, does not change markedly with radius in NGC 2403. The Hudon et al. Field 1 LF falls well below that of the Region I_g and Hudon et al. Field 2 + 3 LFs at $M_{bol} = -4.5$, although this may be due to incompleteness (see text). The errorbar at the faint end of the Field 1 LF shows the uncertainties listed in Table V of Hudon et al. (1989).









































